

# AP Chemistry Equilibrium Worksheet

apchemequiws0809.doc

Name: \_\_\_\_\_


Date: \_\_\_\_\_ Period: \_\_\_\_\_

1. Carbon monoxide and hydrogen react according to the following equation:




When 1.000mol CO and 3.000mol H<sub>2</sub> are placed in a 10.00L vessel at 927°C (1200.K) and allowed to come to equilibrium, the mixture is found to contain 0.387mol H<sub>2</sub>O. What is the molar composition of the equilibrium mixture? That is, how many moles of each substance are present?


2. a. Write the equilibrium constant expression K<sub>c</sub> for catalytic methanation.




b. Write the equilibrium constant expression K<sub>c</sub> for the reverse of the previous reaction




c. Write the equilibrium constant expression K<sub>c</sub> for the synthesis of ammonia.



d. Write the equilibrium expression K<sub>c</sub> when the equation for the previous reaction is written



3. Hydrogen iodide, HI, decomposes at moderate temperatures according to the equation



The amount of I<sub>2</sub> in the reaction mixture can be determined from the intensity of the violet color of I<sub>2</sub>; the more intense the color, the more I<sub>2</sub> in the reaction vessel. When 4.00mol HI was placed in a 5.00L vessel at 458°C, the equilibrium mixture was found to contain 0.442mol I<sub>2</sub>. What is the value of K<sub>c</sub> for the decomposition of HI at this temperature?

4. Phosphorus pentachloride dissociates on heating:



If K<sub>c</sub> equals 3.28x10<sup>-2</sup> at 191°C, what is K<sub>p</sub> at this temperature?

5. a. Quicklime (calcium oxide, CaO) is prepared by heating a source of calcium carbonate, CaCO<sub>3</sub>, such as limestone or seashells.




Write the expression for K<sub>c</sub>.


b. You can write the equilibrium constant expression for a physical equilibrium, such as vaporization, as well as for a chemical equilibrium. Write the expression for K<sub>c</sub> for the vaporization of water.



6. A 50.0L reaction vessel contains 1.00mol N<sub>2</sub>, 3.00mol H<sub>2</sub>, and 0.500mol NH<sub>3</sub>. Will more ammonia, NH<sub>3</sub>, be formed or will it dissociate when the mixture goes to equilibrium at 400.°C? The equation is:




7. A gaseous mixture contains 0.30mol CO, 0.10mol H<sub>2</sub>, and 0.020mol H<sub>2</sub>O, plus an unknown amount of CH<sub>4</sub>, in each liter. This mixture is at equilibrium at 1200.K.




What is the concentration of CH<sub>4</sub> in this mixture? The equilibrium constant K<sub>c</sub> equals 3.92.

8. The reaction




is used to increase the ratio of hydrogen in synthesis gas (mixtures of CO and H<sub>2</sub>). Suppose you start with 1.00mol each of carbon monoxide and water in a 50.0L vessel. How many moles of each substance are in the equilibrium mixture at 1000.°C? The equilibrium constant K<sub>c</sub> at this temperature is 0.58.

9. Hydrogen and iodine react according to the equation






Suppose 1.00mol H<sub>2</sub> and 2.00mol I<sub>2</sub> are placed in a 1.00L vessel. How many moles of substances are in the gaseous mixture when it comes to equilibrium at 458°C? The equilibrium constant K<sub>c</sub> at this temperature is 49.7.


10. Predict the direction of reaction when H<sub>2</sub> is removed from a mixture (lowering its concentration) in which the following equilibrium has been established:



11. Look at each of the following equations and decide whether an increase of pressure obtained by decreasing the volume will increase, decrease, or have no effect on the amounts of products.



12. Carbon monoxide is formed when carbon dioxide reacts with solid carbon (graphite).



Is a high or low temperature more favorable to the formation of carbon monoxide?